


INTERACTION BETWEEN SURFACE AND INTERNAL WAVES IN SHALLOW WATER

Analysis of data acquired at the U.S. Navy Electronics Laboratory
Oceanographic Research Tower site

W. Krauss • Research and Development Report • 30 January 1967
U. S. NAVY ELECTRONICS LABORATORY, SAN DIEGO, CALIFORNIA

TR
7855
105
no. 1432

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED



PROBLEM

Investigate and report on the nature of internal waves of 2-to-20-minute periods at the U. S. Navy Electronics Laboratory (NEL) Oceanographic Research Tower. Specifically, study their relation to surface waves.

RESULTS

1. Resonant interaction between amplitude-modulated swell and internal waves may create internal waves of 2-to-20-minute periods. Internal waves with amplitudes of 1-3 meters can be produced within an interaction time of 15 minutes if the stratification of the water is such that

$$|\vec{k}_1 - \vec{k}_2|^2 / (\omega_1 - \omega_2)^2$$

is an eigenvalue of the internal wave equation (\vec{k}_1 , \vec{k}_2 , ω_1 , and ω_2 being wave numbers and frequencies of the modulated swell).

2. Internal waves due to amplitude-modulated swell have the same characteristics as the modulation. Specifically, they have the same wavelength and period and travel in the same direction as the amplitude modulation.

3. The resonance process is most efficient in the case of a modulation which travels in the same direction as the carrier wave (the main constituent) of a swell. The creation of internal waves of this type is strongly dependent on the stratification of the water.

4. Preliminary evaluations of temperature and swell records from the NEL Tower show good agreement between periods of swell modulation and internal wave periods.

5. It is likely that internal waves are created on the entire continental shelf off Southern California during times of favorable stratification. This depends on tides and wind.

RECOMMENDATIONS

1. Verify, with additional measurements at the NEL Tower, the theory that internal waves are produced by swell.
2. Acquire, for complete analysis, long records of temperature fluctuations and surface waves in order to compute spectra with high resolution and statistical confidence.



3. Establish, by measurements of surface waves at two positions, and slick observations, that internal waves travel in the direction of the swell modulation.

ADMINISTRATIVE INFORMATION

Work was performed under SR 104 03 01, Task 0594 (NEL L41171). The report covers work from August 1966 to December 1966, and was approved for publication 30 January 1967. Financial arrangements for travel and salary were furnished by the Office of Naval Research (ONR) and administered by Scripps Institution of Oceanography.

The author is grateful for the assistance of J. L. Cairns, D. E. Good, J. R. Olson, and C. L. Barker in collecting data; of Mrs. R. P. Brown in programming and computer analysis; of C. V. Hart in computations and plotting; and E. C. LaFond, O. S. Lee, and J. L. Cairns for discussions.

CONTENTS

LIST OF SYMBOLS . . .	page 5
INTRODUCTION . . .	7
ENERGY TRANSFER FROM SURFACE WAVES TO INTERNAL WAVES BY RESONANCE . . .	9
EQUATIONS AND BOUNDARY CONDITIONS . . .	10
THE PERTURBATION EQUATIONS . . .	11
THE EQUATION FOR $\vec{w}(x, z, t)$. . .	13
THE PRIMARY WAVE FIELD AND THE FORCES f_1 and f_2 . . .	15
THE SECONDARY WAVE FIELD IN THE RESONANCE CASE . . .	17
$F_{mn}^{\pm}(z)$, $G_{mn}^{\pm}(0)$, and A_{mnr}^{\pm} IN CASE OF AN EXPONENTIALLY STRATIFIED SEA . . .	19
THE SECONDARY RESONANCE WAVE FIELD DUE TO SWELL IN CASE OF AN EXPONENTIALLY STRATIFIED SEA . . .	23
MEASUREMENTS ON SWELL AND INTERNAL WAVES . . .	24
CONCLUSIONS . . .	26
RECOMMENDATIONS . . .	27
REFERENCES . . .	27

ILLUSTRATIONS

- 1 Temperature recorded by thermistors 3 to 21 at the NEL Tower (4 October 1966, 1900-2000) . . . page 8
- 2 Amplitude (cm) of internal waves due to modulated swell after an interaction time of 10^3 seconds (16.6 minutes) as a function of the angle between the two primary waves . . . 23
- 3 Swell spectra of wave height sensors 1 and 2 at positions near the NEL Tower, and phase difference between both records (4 October 1966, 1230-1330, sampling rate 1 second); the arrow indicates the 95-percent confidence limit . . . 25
- 4 Histograms of periods of swell modulations (left) and of internal waves (right) at the NEL Tower . . . 26

LIST OF SYMBOLS

If not defined in the article, the symbols have the following meaning:

A_n	Amplitude of W_n
A_{mnr}^{\pm}	Expansion coefficients, see equations (58) and (62)
a_m, a_n	See equations (63) and (64)
F_{mn}^{\pm}	See equation (57)
f_1, f_2	See equations (39) and (40)
G_{mn}^{\pm}	See equation (52)
g	Acceleration of gravity
H	Bottom depth
h	Horizontal components, used as an index
$\vec{k} \equiv (\kappa_1 \eta)$	Wave number vector
p	Pressure, \bar{p} mean pressure
\vec{u}	(u, v, w) orbital current velocity of the waves, with components u, v, w in direction x, y, z
$W_n(z)$	Eigenfunction of n^{th} mode, according to equations (46) to (48)
\vec{x}	(x, y) , horizontal plane
z	Vertical coordinate, pointing downward
$\Gamma(z)$	$\frac{1}{\bar{\rho}} \frac{d\bar{\rho}}{dz}$; $\Gamma_0 = \text{const}$
ζ	Amplitude of a wave
ρ	Density, $\bar{\rho}(z)$ mean density distribution
ϕ	Gravitational potential
ω	Angular frequency
∇	$\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right)$, del operator
$\nabla \cdot$	Divergence
∇^2	Laplace operator

INTRODUCTION

It was shown by E. C. LaFond¹⁻³ and O. S. Lee⁴ that internal waves of 2-to-20-minute periods are a dominant feature in the thermal structure of the sea around the NEL Oceanographic Research Tower. From extensive measurements, E. C. LaFond derived that 50 percent of all waves had periods greater than 7.3 minutes and 50 percent of waves had heights of more than 5.6 feet (170 centimeters). Wave heights of more than 20 feet (6 meters) can sometimes be observed in water only 60 feet (18 meters) deep.

These short-period oscillations are only one part of the entire internal wave spectrum. They are superimposed on longer fluctuations of the mean thermocline, with changes mainly due to internal tides and wind.⁵ Nevertheless, these waves in the 2-to-20-minute range are the most striking fluctuations besides the internal tides. Their occurrence seems to depend on several factors. There is neither a close relationship to the surface tides nor to the internal tides, but the changing stratification due to the tides seems to be of importance.

Most likely, these waves are long-crested, progressive waves traveling toward shore with a velocity of 20-to-40 feet per minute (10-to-20 centimeters per second). The measurements by O. S. Lee⁴ indicate a beamwidth of only ± 15 degrees. This agrees with former measurements by C. W. Ufford,⁶ G. Ewing,⁷ and E. C. LaFond.¹ C. S. Cox⁸ got similar results. They are supported by observations of sea surface slicks, which are often closely related to internal waves. According to O. S. Lee,⁴ the mean speed is 27 feet per minute (13.7 centimeters per second) and the direction 85 degrees.

Figure 1 gives an example of these waves measured on 4 October 1966, 1900-2000. The temperature fluctuations are shown for thermistors 3 to 21. The distance between the thermistors is 2.5 feet, thermistor 3 being 5 feet above the bottom and thermistor 21 about 20 feet below sea surface. The water depth is 60 feet.

After a calm period of several hours the waves start to occur at about 1900. The isotherm depth decreases during the next half hour and during that time high-amplitude internal waves are present. The period is not quite independent of depth. Shorter periods are generally observed near the surface than in deeper layers, but all waves are of first mode. This seems typical for the area.

The origin of these waves is rather obscure. There are no obvious meteorological or tidal forces that could produce the regular wave trains. If there were a constant coupling between the tidal phase and the occurrence of these waves, one would be inclined to interpret them as an adaptation of the changing mean stratification. But this is not possible. The only reason for their creation therefore seems to be surface waves.

F. K. Ball⁹ has shown that, in the case of a two-layered model, resonance is possible for second order interactions between surface and internal boundary waves. S. A. Thorpe¹⁰ extended the theory to wave interactions in a continuously stratified fluid. He showed that a transfer of energy from surface to internal waves may occur, and an internal wave generation mechanism will exist. The theory has been applied to situations which might be realized in the laboratory, but an application to natural conditions has not been attempted. For tank experiments (under somewhat extreme conditions) he found that the internal wave amplitude will be equal to that of the surface waves after an interaction time of only 28 seconds.




Figure 1. Temperature recorded by thermistors 3 to 21 at the NEL Tower (4 October 1966, 1900-2000). (Each curve represents the variation of the temperature from the mean for that numbered thermistor.)

K. E. Kenyon¹¹ extended the theory to a description of the entire internal wave spectrum which may result from the interaction of surface waves in the ocean. He gave a computation of scattering of swell energy into internal wave energy, using observations from the NEL Tower, from which he concluded that this process may be of minor importance for the creation of internal waves.

The following computations are similar to his, but we do not compute the energy spectra, and derive only the equations for the amplitudes of the waves. We come to different results, the reason for this being that Kenyon considered surface wave spectra only for the case that the directionality of the spectrum is given by $\cos^4\alpha$. This means that all surface waves are contained within an angle of about ± 30 degrees. From this narrow directional swell spectrum, energy is fed to internal waves traveling mainly perpendicular to the swell. This, obviously, does not agree with the observations mentioned above. Both swell and internal waves travel mainly toward shore in the sea off Southern California.

Kenyon's conclusion that surface waves with a narrow directional spectrum do not feed considerable energy into the internal wave spectrum is supported by K. Hasselmann.¹² This agrees with our results, but it will be shown here that the process is about 100 times more effective for amplitude-modulated swell, which may be described by two surface waves traveling in directions which include an angle between 90 and 180 degrees. This still leads to a progressive wave, if the two amplitudes differ considerably in magnitude. If these two waves are given by

$$\zeta_1(\vec{x}, t) = A_1 \cos(\vec{k}_1 \cdot \vec{x} - \omega_1 t) \quad , \quad \zeta_2(\vec{x}, t) = A_2 \cos(\vec{k}_2 \cdot \vec{x} - \omega_2 t) \quad (1)$$

with $A_1 \gg A_2$, the resulting wave can be described by

$$\zeta(\vec{x}, t) = \zeta_1(\vec{x}, t) + \zeta_2(\vec{x}, t) = A(\vec{x}, t) \cos(\vec{k}_1 \cdot \vec{x} - \omega_1 t - \phi(\vec{x}, t)) \quad (2)$$

where the amplitude $A(\vec{x}, t)$ and the phase $\phi(\vec{x}, t)$ are given according to

$$A(\vec{x}, t) = \left[A_1^2 + A_2^2 + 2A_1A_2 \cos\left[(\vec{k}_1 - \vec{k}_2) \cdot \vec{x} - (\omega_1 - \omega_2)t\right] \right]^{1/2} \quad (3)$$

and

$$\phi(\vec{x}, t) = \arctan \frac{A_2 \sin\left[(\vec{k}_1 - \vec{k}_2) \cdot \vec{x} - (\omega_1 - \omega_2)t\right]}{A_1 + A_2 \cos\left[(\vec{k}_1 - \vec{k}_2) \cdot \vec{x} - (\omega_1 - \omega_2)t\right]} \quad (4)$$

thus representing a swell which travels in a direction given by the wave vector \vec{k}_1 with an amplitude changing between $A_1 \pm A_2$ and a variable phase. Wave measurements at the NEL Tower indicate the reality of such a swell.

ENERGY TRANSFER FROM SURFACE WAVES TO INTERNAL WAVES BY RESONANCE

The process of resonant interaction is easy to comprehend, but the detailed analysis involves a great deal of algebra. A short outline of the main ideas may help in understanding the following computations.

The hydrodynamic equations are nonlinear. They include terms of the form $\vec{u} \cdot \nabla \vec{u}$, \vec{u} being the velocity vector. In internal wave theory, these equations are linearized by perturbation methods. The result is a sum of mutually independent waves, each one conserving its own energy if friction is neglected. Many features are adequately described by this linear theory, but there exist circumstances under which the nonlinear terms can give rise to significant energy transfer between these primary wave solutions.

Surface waves are zero mode solutions of the internal wave equation. Internal waves may be of any mode larger than zero.

The computations in the following sections are based on this idea. Consider the simple nonlinear wave equation

$$\frac{\partial^2 \phi}{\partial t^2} + \omega^2 \phi - \phi^2 = 0 \quad (5)$$

Applying perturbation methods, $\phi = \phi^{(1)} + \phi^{(2)} + \dots$, the first-order equation reads

$$\frac{\partial^2 \phi^{(1)}}{\partial t^2} + \omega^2 \phi^{(1)} = 0 \quad (6)$$

with the solution

$$\phi^{(1)} = \sum_n a_n e^{i\omega_n t} \quad (7)$$

These are mutually independent waves. The second-order equation is

$$\frac{\partial^2 \phi^{(2)}}{\partial t^2} + \omega^2 \phi^{(2)} = \phi^{(1)2} = \sum_m \sum_n a_m a_n e^{i(\omega_m + \omega_n)t} \quad (8)$$

The secondary waves $\phi^{(2)}$ are therefore forced waves with frequencies $\omega_m + \omega_n$. As long as these combination frequencies $\omega_m + \omega_n$ are not the same as a natural frequency of the system, the amplitudes of the secondary waves will remain small. However, when one of the combination frequencies $\omega_m + \omega_n$ is the same as one of the natural frequencies ω of the system, this wave never gets out of phase with the forcing wave, and a continuous energy transfer, only limited by the available energy in the primary waves, is possible. If this occurs, the solution of (8) is of the type $\phi^{(2)} \propto t$, that is, $\phi^{(2)}$ increases linearly with time. This is the resonance case. Its physical meaning is that energy from frequencies ω_n and ω_m is continuously fed into the frequency range $\omega_m + \omega_n$. We will study the problem of whether energy from surface waves in the frequency bands ω_m and ω_n can be transferred into internal waves in the frequency band $\omega_m \pm \omega_n$. Obviously, if ω_m and ω_n are swell frequencies, a transfer to the frequency band $\omega_m - \omega_n$ is possible only because internal waves have a cutoff at the Väisälä frequency, which is considerably lower than the swell frequencies. Additionally, the frequencies ω_m and ω_n must be very close together in order to feed energy into a frequency band corresponding to periods 2 to 20 minutes.

EQUATIONS AND BOUNDARY CONDITIONS

The hydrodynamic equations of a nonviscous, incompressible, stably stratified fluid in a Cartesian frame of reference with z pointing downward may be written as

$$\rho \frac{\partial \vec{u}}{\partial t} + \rho \vec{u} \cdot \nabla \vec{u} = -\nabla p - \rho \nabla \phi \quad (9)$$

$$\frac{\partial \rho}{\partial t} + \vec{u} \cdot \nabla \rho = 0 \quad (10)$$

$$\nabla \cdot \vec{u} = 0 \quad (11)$$

The symbols have the following meaning: $\vec{u} \equiv (u, v, w)$ is the fluid velocity, p the pressure, ϕ the gravitational potential, and ρ the density. The effect of the earth's rotation is neglected. The boundary conditions are

$$w = -\left(\frac{\partial \zeta}{\partial t} + \vec{u} \cdot \nabla \zeta\right) \text{ at the sea surface, } z = -\zeta(\vec{x}, t) \quad (12)$$

$$p = \text{const at the sea surface, } z = -\zeta(\vec{x}, t) \quad (13)$$

$$w = 0 \text{ at the bottom, } z = H = \text{const} \quad (14)$$

In order to apply the boundary conditions (12) and (13) at the undisturbed sea surface $z = 0$, we expand them about $z = 0$. The Taylor series of (12) is

$$w - \zeta \frac{\partial w}{\partial z} + \frac{\zeta^2}{2} \frac{\partial^2 w}{\partial z^2} + \dots = -\left[\frac{\partial \zeta}{\partial t} + \left(\vec{u} - \zeta \frac{\partial \vec{u}}{\partial z} + \frac{\zeta^2}{2} \frac{\partial^2 \vec{u}}{\partial z^2} + \dots\right) \cdot \nabla \zeta\right] \text{ at } z=0 \quad (15)$$

equation (13) yields

$$p - \zeta \frac{\partial p}{\partial z} + \frac{\zeta^2}{2} \frac{\partial^2 p}{\partial z^2} + \dots = \text{const at } z = 0 \quad (16)$$

THE PERTURBATION EQUATIONS

Suppose the variables in (9) to (11) can be written as perturbation series

$$\begin{aligned} \vec{u} &= \vec{u}^{(1)} + \vec{u}^{(2)} + \dots, \quad p = p^{(0)} + p^{(1)} + p^{(2)} + \dots, \quad \rho = \rho^{(0)} + \rho^{(1)} + \rho^{(2)} + \dots, \\ \zeta &= \zeta^{(1)} + \zeta^{(2)} + \dots, \end{aligned} \quad (17)$$

with $p^{(0)} = \bar{p}(z)$ and $\rho^{(0)} = \bar{\rho}(z)$, then, entering the equations (9) to (11) and (15) to (16), we arrive at the following equations, omitting terms of higher than second order.

0th Order Equations

$$\frac{d\bar{p}}{dz} = g\bar{\rho} \quad (18)$$

with

$$\bar{p} = \text{const at } z = 0. \quad (19)$$

1st Order Equations

$$\bar{\rho} \frac{\partial \vec{u}^{(1)}}{\partial t} + \nabla p^{(1)} + \rho^{(1)} \nabla \phi = 0 \quad (20)$$

$$\frac{\partial \rho^{(1)}}{\partial t} + w^{(1)} \frac{d\bar{\rho}}{dz} = 0 \quad (21)$$

$$\nabla \cdot \vec{u}^{(1)} = 0 \quad (22)$$

with

$$w^{(1)} = - \frac{\partial \zeta^{(1)}}{\partial t} \text{ at } z = 0 \quad (23)$$

$$p^{(1)} - \zeta^{(1)} \frac{d\bar{p}}{dz} = 0 \text{ at } z = 0 \quad (24)$$

$$w^{(1)} = 0 \text{ at } z = H \quad (25)$$

2nd Order Equations

$$\bar{\rho} \frac{\partial \vec{u}^{(2)}}{\partial t} + \nabla p^{(2)} + \rho^{(2)} \nabla \phi = - \left(\rho^{(1)} \frac{\partial \vec{u}^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla \vec{u}^{(1)} \right) \quad (26)$$

$$\frac{\partial \rho^{(2)}}{\partial t} + w^{(2)} \frac{d\bar{\rho}}{dz} = - \vec{u}^{(1)} \cdot \nabla \rho^{(1)} \quad (27)$$

$$\nabla \cdot \vec{u}^{(2)} = 0 \quad (28)$$

with

$$w^{(2)} + \frac{\partial \zeta^{(2)}}{\partial t} = \zeta^{(1)} \frac{\partial w^{(1)}}{\partial z} - \vec{u}^{(1)} \cdot \nabla \zeta^{(1)} \text{ at } z = 0 \quad (29)$$

$$p^{(2)} - \zeta^{(2)} \frac{d\bar{p}}{dz} = \zeta^{(1)} \frac{\partial p^{(1)}}{\partial z} - \frac{\zeta^{(1)2}}{2} \frac{d^2 \bar{p}}{dz^2} \text{ at } z = 0 \quad (30)$$

$$w^{(2)} = 0 \text{ at } z = H \quad (31)$$

The 0th order equations characterize the mean stable conditions; the 1st order ones describe linear internal and surface waves in that fluid; and the 2nd order equations give nonlinear effects of these waves, including energy transfers to other frequency bands. It is this set of equations which leads to energy flux from surface waves to internal waves.

THE EQUATIONS FOR $w(\vec{x}, z, t)$

We intend to eliminate u , v , ρ , and p in the left-hand sides of the 1st and 2nd order equations in order to get equations for w . The procedure is well-known.¹³ For the 1st order equations we get

$$\nabla_h \cdot \left(\frac{\partial^2 w^{(1)}}{\partial t^2} + g \Gamma w^{(1)} \right) + \frac{\partial^4 w^{(1)}}{\partial t^2 \partial z^2} + \Gamma \frac{\partial^3 w^{(1)}}{\partial t^2 \partial z} = 0 \quad (32)$$

where

$$\Gamma(z) = \frac{1}{\bar{\rho}} \frac{d\bar{\rho}}{dz}$$

Combining the boundary conditions (23) and (24) together with the horizontal components of (20) and the continuity equation (22), we find

$$\frac{\partial^3 w^{(1)}}{\partial t^2 \partial z} + g \nabla_h^2 w^{(1)} = 0 \text{ at } z = 0 \quad (33)$$

and (25) remains the same

$$w^{(1)} = 0 \text{ at } z = H \quad (34)$$

Internal and surface waves of the first order are governed by equation (32) with boundary conditions (33) and (34). The problem has been solved analytically

for many density distributions $\bar{\rho}(z)$ and may be handled numerically for any distribution $\bar{\rho}(z)$. The result is a surface wave $w_0^{(1)}$ and a set of internal waves

$$\sum_{n=1}^{\infty} w_n^{(1)}, \text{ each one being mutually independent and traveling with a phase}$$

velocity given by the eigenvalues of the problem.

The same procedure is used for deriving an equation for $w^{(2)}$ from (26) to (28): The operator $\nabla_h \cdot$ applied to (26) together with (28) yields

$$-\bar{\rho} \frac{\partial^2 w^{(2)}}{\partial t \partial z} + \nabla_h^2 p^{(2)} = -\nabla_h \cdot \left(\rho^{(1)} \frac{\partial \vec{u}_h^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla \vec{u}_h^{(1)} \right) \quad (35)$$

Differentiating with respect to z and t we have

$$-\frac{d\bar{\rho}}{dz} \frac{\partial^3 w^{(2)}}{\partial t^2 \partial z} - \bar{\rho} \frac{\partial^4 w^{(2)}}{\partial t^2 \partial z^2} + \nabla_h^2 \frac{\partial^2 p^{(2)}}{\partial z \partial t} = -\frac{\partial^2}{\partial z \partial t} \nabla_h \cdot \left(\rho^{(1)} \frac{\partial \vec{u}_h^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla \vec{u}_h^{(1)} \right)$$

Combining the vertical component of (26) with (27) gives

$$\bar{\rho} \frac{\partial^2 w^{(2)}}{\partial t^2} + \frac{\partial^2 p^{(2)}}{\partial z \partial t} + g \frac{d\bar{\rho}}{dz} w^{(2)} = -g \vec{u}^{(1)} \cdot \nabla \rho^{(1)} - \frac{\partial}{\partial t} \left(\rho^{(1)} \frac{\partial w^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla w^{(1)} \right),$$

which after applying ∇_h^2 and substituting $p^{(2)}$ in both equations above yields

$$\begin{aligned} \nabla_h^2 \left(\frac{\partial^2 w^{(2)}}{\partial t^2} + g \Gamma w^{(2)} \right) + \frac{\partial^4 w^{(2)}}{\partial t^2 \partial z^2} + \Gamma \frac{\partial^3 w^{(2)}}{\partial t^2 \partial z} = & -\frac{1}{\bar{\rho}} \nabla_h^2 \left[g \vec{u}^{(1)} \cdot \nabla \rho^{(1)} \right. \\ & \left. + \frac{\partial}{\partial t} \left(\rho^{(1)} \frac{\partial w^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla w^{(1)} \right) \right] + \frac{1}{\bar{\rho}} \frac{\partial^2}{\partial z \partial t} \nabla_h \cdot \left(\rho^{(1)} \frac{\partial \vec{u}_h^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla \vec{u}_h^{(1)} \right) \end{aligned} \quad (36)$$

Combining the boundary conditions (29) and (23) together with (35) and eliminating $\frac{\partial p^{(1)}}{\partial z}$ in the inhomogeneous part by means of the vertical component of (26), we find

$$\begin{aligned} \frac{\partial^3 w^{(2)}}{\partial t^2 \partial z} + g \nabla_h^2 w^{(2)} = & \frac{\partial}{\partial t} \nabla_h \cdot \left(\frac{\rho^{(1)}}{\bar{\rho}} \frac{\partial \vec{u}^{(1)}}{\partial t} + \vec{u}^{(1)} \cdot \nabla \vec{u}_h^{(1)} \right) + \frac{\partial}{\partial t} \nabla_h^2 \left[\left(\frac{g \rho^{(1)}}{\bar{\rho}} - \frac{\partial w^{(1)}}{\partial t} \right) \zeta^{(1)} \right. \\ & \left. - \frac{g \Gamma \zeta^{(1)2}}{2} \right] + g \nabla_h^2 \left(\zeta^{(1)} \frac{\partial w^{(1)}}{\partial z} - \vec{u}_h^{(1)} \cdot \nabla_h \zeta^{(1)} \right) \text{ at } z = 0 \end{aligned} \quad (37)$$

and (31) remains unaltered

$$w^{(2)} = 0 \text{ at } z = H \quad (38)$$

There are two driving forces responsible for second order internal waves:

- (i) The body force f_1 in the fluid due to the nonlinear terms in the equations of motion and the incompressibility condition. It is given by the inhomogeneous part of (36)

$$f_1(\vec{x}, z, t) = -\frac{1}{\bar{\rho}} \nabla_h^2 \left\{ g\vec{u}^{(1)} \cdot \nabla \rho^{(1)} + \frac{\partial}{\partial t} \left(\rho^{(1)} \frac{\partial w^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla w^{(1)} \right) \right\} \\ + \frac{1}{\bar{\rho}} \frac{\partial^2}{\partial z \partial t} \nabla_h \cdot \left(\rho^{(1)} \frac{\partial \vec{u}_h^{(1)}}{\partial t} + \bar{\rho} \vec{u}^{(1)} \cdot \nabla \vec{u}_h \right) \quad (39)$$

- (ii) The surface force f_2 due to the deviations of the sea surface from its mean level, given by the inhomogeneous part of (37)

$$f_2(\vec{x}, z=0, t) = \frac{\partial}{\partial t} \nabla_h \cdot \left(\frac{\rho^{(1)}}{\bar{\rho}} \frac{\partial \vec{u}_h^{(1)}}{\partial t} + \vec{u}^{(1)} \cdot \nabla \vec{u}_h^{(1)} \right) + \frac{\partial}{\partial t} \nabla_h^2 \left[\left(\frac{g\rho^{(1)}}{\bar{\rho}} - \frac{\partial w^{(1)}}{\partial t} \right) \zeta^{(1)} \right. \\ \left. - \frac{g\Gamma\zeta^{(1)2}}{2} \right] + g \nabla_h^2 \left(\zeta^{(1)} \frac{\partial w^{(1)}}{\partial z} - \vec{u}_h^{(1)} \cdot \nabla_h \zeta^{(1)} \right) \quad (40)$$

Comparing these equations with the simple example given at the beginning, the problem (9) to (14) corresponds to equation (5), and (32) to (34) corresponds to (6), and (36) to (38) corresponds to (8).

THE PRIMARY WAVE FIELD AND THE FORCES f_1 AND f_2

Suppose that the primary waves are progressive waves traveling in a direction given by the wave number vector $\vec{k} \equiv (\kappa, \eta)$. We will use the notation $k \equiv |\vec{k}| = \sqrt{\kappa^2 + \eta^2}$. A set of solutions of (32) to (34) is

$$\vec{u}_h^{(1)}(\vec{x}, z, t) = - \sum_{n=1}^{\infty} A_n \frac{\vec{k}_n}{k_n^2} \frac{dW_n}{dz} \sin(\vec{k}_n \cdot \vec{x} - \omega_n t) \quad (41)$$

$$w^{(1)}(\vec{x}, z, t) = \sum_{n=1}^{\infty} A_n W_n \cos(\vec{k}_n \cdot \vec{x} - \omega_n t) \quad (42)$$

$$\rho^{(1)}(\vec{x}, z, t) = \frac{d\bar{\rho}}{dz} \sum_{n=1}^{\infty} \frac{A_n}{\omega_n} W_n \sin(\vec{k}_n \cdot \vec{x} - \omega_n t) \quad (43)$$

$$p^{(1)}(\vec{x}, z, t) = -\bar{\rho} \sum_{n=1}^{\infty} A_n \frac{\omega_n}{k_n^2} \frac{dW_n}{dz} \sin(\vec{k}_n \cdot \vec{x} - \omega_n t) \quad (44)$$

$$\zeta^{(1)}(\vec{x}, z, t) = -\frac{1}{g} \sum_{n=1}^{\infty} A_n \frac{\omega_n}{k_n^2} \frac{dW_n}{dz} \Big|_{z=0} \sin(\vec{k}_n \cdot \vec{x} - \omega_n t) \quad (45)$$

where $W_n(z)$ is governed by

$$\frac{d^2 W_n}{dz^2} + \Gamma \frac{dW_n}{dz} + (g\Gamma - \omega_n^2) \frac{k_n^2}{\omega_n^2} W_n = 0 \quad (46)$$

with

$$\frac{dW_n}{dz} + \frac{gk_n^2}{\omega_n^2} W_n = 0 \quad \text{at } z = 0 \quad (47)$$

and

$$W_n = 0 \quad \text{at } z = H \quad (48)$$

and represents either surface or internal waves. From (41) to (45) we can derive the driving functions f_1 and f_2 according to (39) and (40). This includes a great deal of algebra. The final result is

$$f_1(\vec{x}, z, t) = -\frac{1}{2} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left\{ F_{mn}^+(z) \sin\left[(\vec{k}_m + \vec{k}_n) \cdot \vec{x} - (\omega_m + \omega_n)t\right] \right. \\ \left. + F_{mn}^-(z) \sin\left[(\vec{k}_m - \vec{k}_n) \cdot \vec{x} - (\omega_m - \omega_n)t\right] \right\} \quad (49)$$

$$f_2(\vec{x}, 0, t) = \frac{1}{2} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left\{ G_{mn}^+(0) \sin\left[(\vec{k}_m + \vec{k}_n) \cdot \vec{x} - (\omega_m + \omega_n)t\right] \right. \\ \left. + G_{mn}^-(0) \sin\left[(\vec{k}_m - \vec{k}_n) \cdot \vec{x} - (\omega_m - \omega_n)t\right] \right\} \quad (50)$$

where $F_{mn}^{\pm}(z)$ and $G_{mn}^{\pm}(0)$ are given by

$$F_{mn}^{\pm}(z) = \left\{ \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2 \omega_n} \left[g\Gamma \pm \omega_n(\omega_m \pm \omega_n) \right] \frac{dW_m}{dz} W_n \mp \left[g\Gamma \pm \omega_n(\omega_m \pm \omega_n) \right] \frac{1}{\omega_n} W_m \frac{dW_n}{dz} \right. \\ \mp \left[\frac{g}{\omega_n} \frac{1}{\bar{\rho}} \frac{d^2 \bar{\rho}}{dz^2} - \frac{\omega_n(\omega_m \pm \omega_n)}{\omega_m} \Gamma \right] W_m W_n \left. \right\} |\vec{k}_m \pm \vec{k}_n|^2 \\ + \left\{ \mp \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \frac{1}{\bar{\rho}} \frac{d}{dz} \left(\bar{\rho} \frac{dW_m}{dz} \frac{dW_n}{dz} \right) \right.$$

$$\begin{aligned}
& + \frac{1}{\bar{\rho}} \frac{d}{dz} \left(\bar{\rho} W_m \frac{d^2 W_n}{dz^2} \right) \mp \frac{\omega_n}{\omega_m} \frac{1}{\bar{\rho}} \frac{d}{dz} \left(\frac{d\bar{\rho}}{dz} W_m \frac{dW_n}{dz} \right) \Bigg\} \\
& \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n)
\end{aligned} \tag{51}$$

$$\begin{aligned}
G_{mn}^{\pm}(0) = & \left\{ \left[\pm \frac{\omega_r}{\omega_m} \Gamma W_m \frac{dW_n}{dz} \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \frac{dW_m}{dz} \frac{dW_n}{dz} - W_m \frac{d^2 W_n}{dz^2} \right] \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \right. \\
& \mp \left(\frac{g\Gamma - \omega_m^2}{\omega_m} W_m + \frac{\Gamma}{2} \frac{\omega_m}{k_m^2} \frac{dW_m}{dz} \right) \frac{\omega_n}{gk_m^2} \frac{dW_n}{dz} \left| \vec{k}_m \pm \vec{k}_n \right|^2 (\omega_m \pm \omega_n) \\
& \left. + \left(\frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \omega_n + \omega_m \right) \frac{1}{k_m^2} \frac{dW_m}{dz} \frac{dW_n}{dz} \left| \vec{k}_m \pm \vec{k}_n \right|^2 \right\} \Bigg|_{z=0}
\end{aligned} \tag{52}$$

THE SECONDARY WAVE FIELD IN THE RESONANCE CASE

The secondary wave field is given by (36) to (38), which with (49) and (50) read

$$\begin{aligned}
\nabla_h^2 \left(\frac{\partial^2 w^{(2)}}{\partial t^2} + g \Gamma w^{(2)} \right) + \frac{\partial^4 w^{(2)}}{\partial z^2 \partial t^2} + \Gamma \frac{\partial^3 w^{(2)}}{\partial t^2 \partial z} = & - \frac{1}{2} \sum_m \sum_n F_{mn}^{\pm}(z) \sin \left[(\vec{k}_m \pm \vec{k}_n) \cdot \vec{x} \right. \\
& \left. - (\omega_m \pm \omega_n)t \right]
\end{aligned} \tag{53}$$

$$\frac{\partial^3 w^{(2)}}{\partial t^2 \partial z} + g \nabla_h^2 w^{(2)} = \frac{1}{2} \sum_m \sum_n G_{mn}^{\pm}(0) \sin \left[(\vec{k}_m \pm \vec{k}_n) \cdot \vec{x} - (\omega_m \pm \omega_n)t \right] \tag{54}$$

at $z=0$

$$w^{(2)} = 0 \text{ at } z = H \tag{55}$$

where either the plus or minus sign holds. If eigenvalues of (53) exist, which fulfill the condition

$$\nu_r = \frac{\left| \vec{k}_m \pm \vec{k}_n \right|^2}{(\omega_m \pm \omega_n)^2} \tag{56}$$

the solution of (53) to (55) is

$$w^{(2)} = \sum_{mnr} \left\{ \left[\frac{(\omega_m \pm \omega_n) A_{mnr}^\pm W_r^\pm(z)}{4g |\vec{k}_m \pm \vec{k}_n|^2} + \tilde{W}_r^\pm(z) \right] t \cos \left[(\vec{k}_m \pm \vec{k}_n) \cdot \vec{x} - (\omega_m \pm \omega_n) t \right] \right. \\ \left. + \mathring{W}_r^\pm(z) \sin \left[(\vec{k}_m \pm \vec{k}_n) \cdot \vec{x} - (\omega_m \pm \omega_n) t \right] \right\} \quad (57)$$

where $F_{mn}^\pm(z)$ has been expanded in an eigenfunction series according to

$$\frac{F_{mn}^\pm(z)}{\Gamma(z)} = \sum_r A_{mnr}^\pm W_r^\pm(z) \quad (58)$$

and the functions $W_r^\pm(z)$, $\tilde{W}_r^\pm(z)$, and $\mathring{W}_r^\pm(z)$ are governed by the following equations

$$\left. \begin{aligned} \frac{d^2 W_r^\pm}{dz^2} + \Gamma \frac{dW_r^\pm}{dz} + \frac{[g\Gamma - (\omega_m \pm \omega_n)^2] |\vec{k}_m \pm \vec{k}_n|^2}{(\omega_m \pm \omega_n)^2} W_r^\pm &= 0 \\ \frac{dW_r^\pm}{dz} + \frac{g |\vec{k}_m \pm \vec{k}_n|^2}{(\omega_m \pm \omega_n)^2} W_r^\pm &= 0 \text{ at } z = 0 \\ W_r^\pm &= 0 \text{ at } z = H \end{aligned} \right\} \quad (59)$$

$$\left. \begin{aligned} \frac{d^2 \tilde{W}_r^\pm}{dz^2} + \Gamma \frac{d\tilde{W}_r^\pm}{dz} + \frac{[g\Gamma - (\omega_m \pm \omega_n)^2] |\vec{k}_m \pm \vec{k}_n|^2}{(\omega_m \pm \omega_n)^2} \tilde{W}_r^\pm &= 0 \\ \sum_r \frac{d\tilde{W}_r^\pm}{dz} = \frac{G_{mn}^\pm}{4(\omega_m \pm \omega_n)}, \quad \sum_r \tilde{W}_r^\pm = -\frac{(\omega_m \pm \omega_n) G_{mn}^\pm}{4g |\vec{k}_m \pm \vec{k}_n|^2} &\text{ at } z = 0 \\ \sum_r \tilde{W}_r^\pm &= 0 \text{ at } z = H \end{aligned} \right\} \quad (60)$$

$$\left. \begin{aligned} \frac{d^2 \mathring{W}_r^\pm}{dz^2} + \Gamma \frac{d\mathring{W}_r^\pm}{dz} + \frac{[g\Gamma - (\omega_m \pm \omega_n)^2] |\vec{k}_m \pm \vec{k}_n|^2}{(\omega_m \pm \omega_n)^2} \mathring{W}_r^\pm &= -\frac{2 |\vec{k}_m \pm \vec{k}_n|^2 g \Gamma}{(\omega_m \pm \omega_n)^3} \tilde{W}_r^\pm \\ \frac{d\mathring{W}_r^\pm}{dz} + \frac{g |\vec{k}_m \pm \vec{k}_n|^2}{(\omega_m \pm \omega_n)^2} \mathring{W}_r^\pm &= \frac{A_{mnr}^\pm}{2g |\vec{k}_m \pm \vec{k}_n|^2} \frac{dW_r^\pm}{dz} \text{ at } z = 0 \\ \mathring{W}_r^\pm &= 0 \text{ at } z = H \end{aligned} \right\} \quad (61)$$

The factor $(\omega_m \pm \omega_n) A_{mnr}^{\pm} / (4g |\vec{k}_m \pm \vec{k}_n|^2)$ in (57) is dimensionless; $W_r^{\pm}(z)$ and $\dot{W}_r^{\pm}(z)$ are velocities and $\ddot{W}_r^{\pm}(z)$ is an acceleration.

From (57) to (61) we conclude that the primary waves create new waves of wave number $\vec{k}_m \pm \vec{k}_n$ and frequency $\omega_m \pm \omega_n$ with amplitudes depending on the driving forces $F_{mn}^{\pm}(z)$ and $G_{mn}^{\pm}(0)$. These new waves grow linearly with time if the resonance condition (56) is met. The amplitude factor A_{mnr}^{\pm} is given according to (58) by

$$A_{mnr}^{\pm} = \frac{\int_0^H \frac{[g\Gamma - (\omega_m \pm \omega_n)^2] \bar{\rho}}{\Gamma} F_{mn}^{\pm}(z) W_r^{\pm}(z) dz}{\int_0^H [g\Gamma - (\omega_m \pm \omega_n)^2] \bar{\rho} W_r^{\pm}(z)^2 dz} \quad (62)$$

If more than one internal mode is to be considered, $G_{mn}^{\pm}(0)$ has to be expanded into eigenfunctions, too, and evaluated at $z = 0$. Formulas similar to (58) and (62) will hold then for $G_{mn}^{\pm}(0)$.

$F_{mn}^{\pm}(z)$, $G_{mn}^{\pm}(0)$ AND A_{mnr}^{\pm} IN CASE OF AN EXPONENTIALLY STRATIFIED SEA

In order to evaluate solution (57) of the secondary resonance waves, we have to know the eigenfunctions of $W_r^{\pm}(z)$, which are given by (59). They depend on the mean stratification $\bar{\rho}(z)$ and determine the amplitude factor A_{mnr}^{\pm} , given by (62). Equations (59) can be solved numerically for any stable stratified fluid, and it is known that the eigenvalues may vary considerably due to density changes. However, solving (59) for an exponentially stratified sea, where $\Gamma_0 = \text{const}$, may be sufficient at the present time where only the magnitude of the effects is of interest.

Using $\bar{\rho}(z) = \rho_0 e^{\Gamma_0 z}$ as density distribution, W_n is given as solution of (46) to (48)

$$W_n(z) = A_n e^{-(\Gamma_0/2)z} \sin a_n(z - H) \quad (63)$$

where

$$\tan a_n H = - \frac{a_n}{\frac{\Gamma_0}{2} - \frac{gk_n^2}{\omega_n^2}}, \quad a_n = \frac{1}{2} \sqrt{\frac{4(g\Gamma_0 - \omega_n^2)k_n^2}{\omega_n^2} - \Gamma_0^2} \quad (64)$$

and similar equations hold for $W_m(z)$. Entering equation (51) and (52), we arrive at

$$\begin{aligned}
F_{mn}^{\pm} = & A_m A_n e^{-\Gamma_0 z} \left\{ \left[\frac{|\vec{k}_m \pm \vec{k}_n|^2}{2\omega_n k_m^2} (g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n)) (\mp a_n k_m^2 - a_m \vec{k}_m \cdot \vec{k}_n) \right. \right. \\
& + \frac{a_m - a_n}{2} \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left(a_n^2 - \frac{\Gamma_0^2}{4} \mp \frac{\omega_n}{\omega_m} \frac{\Gamma_0^2}{2} \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \right) \left(a_m a_n \right. \\
& \left. \left. + \frac{\Gamma_0^2}{4} \right) \right] \sin(a_m - a_n)(z - H) + \left[\frac{|\vec{k}_m \pm \vec{k}_n|^2}{2\omega_n k_m^2} (g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n)) (\mp a_n k_m^2 + a_m \vec{k}_m \cdot \vec{k}_n) \right. \\
& + \frac{a_m + a_n}{2} \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left(-a_n^2 + \frac{\Gamma_0^2}{4} \pm \frac{\omega_n}{\omega_m} \frac{\Gamma_0^2}{2} \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \right) \left(a_m a_n \right. \\
& \left. \left. - \frac{\Gamma_0^2}{4} \right) \right] \sin(a_m + a_n)(z - H) + \frac{\Gamma_0}{2} \left[|\vec{k}_m \pm \vec{k}_n|^2 \left[\left(\pm 1 - \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \right) \frac{g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n)}{2\omega_n} \right. \right. \\
& \mp \left(\frac{g\Gamma_0}{\omega_n} - \frac{\omega_n(\omega_m \pm \omega_n)}{\omega_m} \right) \left. \right] + (a_m - a_n) \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left[\mp \frac{\vec{k}_m \cdot \vec{k}_n}{2k_m^2} (a_m - a_n) \right. \\
& + a_n \left(\mp \frac{\omega_n}{\omega_m} - 1 \right) \left. \right] \cos(a_m - a_n)(z - H) + \frac{\Gamma_0}{2} \left[|\vec{k}_m \pm \vec{k}_n|^2 \left[\left(\mp 1 + \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \right) \right. \right. \\
& \left. \left. + \frac{g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n)}{2\omega_n} \pm \left(\frac{g\Gamma_0}{\omega_n} - \frac{\omega_n(\omega_m \pm \omega_n)}{\omega_m} \right) \right] + (a_m + a_n) \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \right. \\
& \left. \left. + \left[\pm \frac{\vec{k}_m \cdot \vec{k}_n}{2k_m^2} (a_m + a_n) + a_n \left(\mp \frac{\omega_n}{\omega_m} - 1 \right) \right] \cos(a_m + a_n)(z - H) \right] \right\} \quad (65)
\end{aligned}$$

$$G_{mn}^{\pm}(0) = \frac{1}{2} A_m A_n (\omega_m \pm \omega_n).$$

$$\begin{aligned}
& \cdot \left\{ - \left[a_n M + \frac{\Gamma_0}{2} (a_m - a_n) N + \frac{\Gamma_0}{2} (1 + a_n) P \right] \sin(a_m - a_n) H \right. \\
& + \left[-a_n M + \frac{\Gamma_0}{2} (a_m - a_n) N - \frac{\Gamma_0}{2} (1 + a_n) P \right] \sin(a_m + a_n) H \\
& + \left[-\frac{\Gamma_0}{2} M + \left(a_m a_n + \frac{\Gamma_0^2}{4} \right) N + \left(a_n^2 - \frac{\Gamma_0^2}{4} \right) P \right] \cos(a_m - a_n) H \\
& \left. + \left[\frac{\Gamma_0}{2} M + \left(a_m a_n - \frac{\Gamma_0^2}{4} \right) N - \left(a_n^2 - \frac{\Gamma_0^2}{4} \right) P \right] \cos(a_m + a_n) H \right\} \quad (66)
\end{aligned}$$

with

$$\begin{aligned}
 M &= \pm \frac{\omega_n \Gamma_0}{\omega_m} \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) \mp \frac{g \Gamma_0 - \omega_m^2}{\omega_m^2} \frac{\omega_n^2}{g k_n^2} \left| \vec{k}_m \pm \vec{k}_n \right|^2 \\
 N &= \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) + \left| \vec{k}_m \pm \vec{k}_n \right|^2 \left[\mp \frac{\Gamma_0}{2} \frac{\omega_m \omega_n}{g k_m^2 k_n^2} + \left(\frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \omega_n \right. \right. \\
 &\quad \left. \left. + \omega_m \right) \frac{1}{k_m^2 (\omega_m \pm \omega_n)} \right] \\
 P &= 1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2}
 \end{aligned}$$

A_{mnr}^\pm follows then from (62) and (65)

$$\begin{aligned}
 A_{mnr}^\pm &= \frac{A_m A_n e^{-(\Gamma_0/2)H}}{\Gamma_0 \left(H - \frac{\sin a_r H \cos a_r H}{a_r} \right)} \\
 &\left\{ \frac{Q^\pm \left[-\frac{\Gamma_0}{2} + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \cos(a_m - a_n - a_r)H + (a_m - a_n - a_r) \sin(a_m - a_n - a_r)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_m - a_n - a_r)^2} \right. \\
 &- \frac{Q^\pm \left[-\frac{\Gamma_0}{2} + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \cos(a_m - a_n + a_r)H + (a_m - a_n + a_r) \sin(a_m - a_n + a_r)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_m - a_n + a_r)^2} \\
 &+ \frac{R^\pm \left[-\frac{\Gamma_0}{2} + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \cos(a_m + a_n - a_r)H + (a_m + a_n - a_r) \sin(a_m + a_n - a_r)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_m + a_n - a_r)^2} \\
 &- \left. \frac{R^\pm \left[-\frac{\Gamma_0}{2} + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \cos(a_m + a_n + a_r)H + (a_m + a_n + a_r) \sin(a_m + a_n + a_r)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_m + a_n + a_r)^2} \right) \\
 &- \frac{S^\pm \frac{\Gamma_0}{2} \left[a_r - a_m + a_n + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \sin(a_r - a_m + a_n)H - (a_r - a_m + a_n) \cos(a_r - a_m + a_n)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_r - a_m + a_n)^2}
 \end{aligned}$$

$$\left. \begin{aligned}
& \frac{S^\pm \frac{\Gamma_0}{2} \left[a_r + a_m - a_n + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \sin(a_r + a_m - a_n)H - (a_r + a_m - a_n) \cos(a_r + a_m - a_n)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_r + a_m - a_n)^2} \\
& \frac{T^\pm \frac{\Gamma_0}{2} \left[a_r - a_m - a_n + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \sin(a_r - a_m - a_n)H - (a_r - a_m - a_n) \cos(a_r - a_m - a_n)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_r - a_m - a_n)^2} \\
& \frac{T^\pm \frac{\Gamma_0}{2} \left[a_r + a_m + a_n + e^{(\Gamma_0/2)H} \left(\frac{\Gamma_0}{2} \sin(a_r + a_m + a_n)H - (a_r + a_m + a_n) \cos(a_r + a_m + a_n)H \right) \right]}{\frac{\Gamma_0^2}{4} + (a_r + a_m + a_n)^2}
\end{aligned} \right\} \quad (67)$$

with

$$\begin{aligned}
Q^\pm &= \frac{|\vec{k}_m \pm \vec{k}_n|^2}{2\omega_n k_m^2} \left[g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n) \right] \left[\mp a_n k_m^2 - a_m \vec{k}_m \cdot \vec{k}_n \right] \\
&\quad + \frac{a_m - a_n}{2} \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left[a_n^2 - \frac{\Gamma_0^2}{4} \mp \frac{\omega_n}{\omega_m} \frac{\Gamma_0^2}{2} \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \left(a_m a_n + \frac{\Gamma_0^2}{4} \right) \right] \\
R^\pm &= \frac{|\vec{k}_m \pm \vec{k}_n|^2}{2\omega_n k_m^2} \left[g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n) \right] \left[\mp a_n k_m^2 + a_m \vec{k}_m \cdot \vec{k}_n \right] \\
&\quad + \frac{a_m + a_n}{2} \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left[-a_n^2 + \frac{\Gamma_0^2}{4} \pm \frac{\omega_n}{\omega_m} \frac{\Gamma_0^2}{2} \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \left(a_m a_n + \frac{\Gamma_0^2}{4} \right) \right] \\
S^\pm &= |\vec{k}_m \pm \vec{k}_n|^2 \left[\left(\pm 1 - \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \right) \frac{g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n)}{2\omega_n} \mp \left(\frac{g\Gamma_0}{\omega_n} - \frac{\omega_n(\omega_m \pm \omega_n)}{\omega_m} \right) \right] \\
&\quad + (a_m - a_n) \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left[\mp \frac{\vec{k}_m \cdot \vec{k}_n}{2k_m^2} (a_m - a_n) + a_n \left(\mp \frac{\omega_n}{\omega_m} - 1 \right) \right] \\
T^\pm &= |\vec{k}_m \pm \vec{k}_n|^2 \left[\left(\mp 1 + \frac{\vec{k}_m \cdot \vec{k}_n}{k_m^2} \right) \frac{g\Gamma_0 \pm \omega_n(\omega_m \pm \omega_n)}{2\omega_n} \pm \left(\frac{g\Gamma_0}{\omega_n} - \frac{\omega_n(\omega_m \pm \omega_n)}{\omega_m} \right) \right] \\
&\quad + (a_m + a_n) \left(1 \pm \frac{\vec{k}_m \cdot \vec{k}_n}{k_n^2} \right) (\omega_m \pm \omega_n) \left[\pm \frac{\vec{k}_m \cdot \vec{k}_n}{2k_m^2} (a_m + a_n) + a_n \left(\mp \frac{\omega_n}{\omega_m} - 1 \right) \right]
\end{aligned} \quad (68)$$

THE SECONDARY RESONANCE WAVE FIELD DUE TO SWELL IN CASE OF AN EXPONENTIALLY STRATIFIED SEA

With reference to observations described in the following section, we evaluated the resonance part of solution (52), using (59), (60), (66), (67), and (68). The results are based on the following numerical values:

The swell frequencies of the two primary waves are $\omega_1 = 3.808 \cdot 10^{-1} \text{ sec}^{-1}$ and $\omega_2 = 3.900 \cdot 10^{-1} \text{ sec}^{-1}$. The corresponding wave numbers for a water depth of $H = 18 \text{ m}$ are $k_1 = 3.00 \cdot 10^{-4} \text{ cm}^{-1}$, $k_2 = 3.08 \cdot 10^{-4} \text{ cm}^{-1}$. The density stratification is given by $\Gamma_0 = 5.811 \cdot 10^{-7} \text{ cm}^{-1}$. The corresponding periods and wavelengths of the primary waves are $\tau_1 = 16.50 \text{ seconds}$, $\tau_2 = 16.11 \text{ seconds}$, $\lambda_1 = 209.4 \text{ m}$, $\lambda_2 = 204.0 \text{ m}$. Furthermore, for surface waves we have $a_n = i|k_n|$, $a_m = i|k_m|$. The product between the amplitudes A_n and A_m of the vertical components of the velocity is supposed to be $A_1 A_2 = 400 \text{ cm}^2 \text{ sec}^{-2}$. The corresponding amplitude product of the two swell waves is $\zeta_1^* \zeta_2^* = 889.76 \text{ cm}^2$, which may be based on $\zeta_1^* \approx 50 \text{ cm}$ and $\zeta_2^* \approx 17.8 \text{ cm}$.

Because of these swell frequencies we expect an internal wave of frequency $\omega_2 - \omega_1 = 0.0092 \text{ sec}^{-1}$ or period $\tau_{2-1} = 11.38 \text{ minutes}$. The result is shown in figure 2, which contains the amplitude of this internal wave after an interaction




Figure 2. Amplitude (cm) of internal waves due to modulated swell after an interaction time of 103 seconds (16.6 minutes) as a function of the angle between the two primary waves.

time of 10^3 sec (≈ 16.7 minutes). The amplitude is given in a logarithmic scale as a function of the angle θ between the two primary swell waves. Amplitudes of about 13 centimeters are obtained if the primary waves are traveling in directions containing an angle of about 60 degrees. The amplitude increases to more than 3 meters for angles of more than 150 degrees, and is only a few centimeters for angles less than 30 degrees. This seems to support the results obtained by K. E. Kenyon¹¹ and K. Hasselmann¹² that a narrow directional spectrum of surface waves does not lead to significant internal waves. On the other hand, the figure demonstrates that surface swell which can be described by two surface waves traveling in different directions may create internal waves very rapidly. A swell of that type would be interpreted according to equations (1) to (4) as amplitude-modulated and traveling in a direction given by \vec{k}_1 , because the amplitude A_1 is much larger than A_2 and therefore would govern the swell.

MEASUREMENTS ON SWELL AND INTERNAL WAVES

In order to test the theory, simultaneous measurements on surface swell and internal waves were made in October 1966 at the NEL Tower, running from 3 October, 1130 to 6 October, 1215. Two wave-height sensors were used to derive the directional spectrum of the swell — one fixed at the NEL Tower, the second at a distance of 39.3 meters toward the west. The voltage output was recorded on punch tape. The temperature fluctuations were measured by means of two vertical thermistor arrays. The spacing between the thermistors was 75 centimeters, reaching from the bottom up to 2.75 meters below mean sea surface. The two arrays were located in positions along a line from southwest to northeast. The distance between the arrays was 284.4 meters. The temperature fluctuations were recorded on an analog recorder and on punch tape. Simultaneous records of swell and temperature fluctuations were obtained. The complete analysis of these data will be published later. Preliminary results support the theory given above. Figure 3 shows the power spectra of the two wave-height sensors for data from 4 October, 1230–1330. The voltage output of the two sensors has been used directly for these calculations. This output is different for both, and this is the reason for spectrum 2 showing higher intensities than spectrum 1. Otherwise, both spectra coincide quite well. The phase difference changes linearly with frequency as is to be expected. But there occurs a remarkable hole at frequencies of about 0.061 cycle per second, corresponding to $\omega \approx 0.38 \text{ sec}^{-1}$ or $\tau \approx 16.4 \text{ sec}$. The energy decreases in both spectra from about 12 to 1.2 (arbitrary units), and the phase differences change from 35° to -70° , which indicates that the swell in this spectral range consists of two waves, one with $\omega_2 = 3.9 \times 10^{-1} \text{ sec}^{-1}$ traveling in direction 30° and a second one with $\omega_1 = 3.808 \times 10^{-1} \text{ sec}^{-1}$ traveling in a direction of about 270° . The angle between both is about 120° . From figure 2 it follows that within a quarter of an hour these two swell waves would create an internal wave of $\tau = 11.4$ minutes with an amplitude of 1.5 meters

The swell in the vicinity of the NEL Tower therefore seems to be modulated as indicated by figure 3. If the stratification is favorable, that is, if $|\vec{k}_m - \vec{k}_a|^2 / (\omega_m - \omega_n)^2$ is an eigenvalue, a very intensive energy transfer from the swell

frequencies ω_m and ω_n to internal waves of the first mode occurs, producing internal waves of amplitudes of several meters. These internal waves have the same characteristics as the swell modulation. Their period coincides with that of the modulation, and the wavelength is the same. Additionally, they travel in the same direction as the modulation.

The agreement between internal wave periods and the periods of the swell modulations was tested. Data from the wave-height sensors for the time from 4 October 1966, 1900 to 5 October 1966, 0145, were used to determine the periods of the modulations. The internal wave periods were taken from the analog records for the time 3 October, 1500 to 5 October, 0300. Histograms of both




Figure 3. Swell spectra of wave height sensors 1 and 2 at positions near the NEL Tower, and phase difference between both records (4 October 1966, 1230-1330; sampling rate, 1 second); the arrow indicates the 95-percent confidence limit.

periods are shown in figure 4. They show the number of occurrences of periods between 2 and 22 minutes in percent. The number of waves contained in each histogram is about 120. The figure is in good agreement with the theory given above, both distributions are peaked in the period range of 4-to-10 minutes and decrease toward periods larger than 20 minutes. Final conclusions, however, are possible only after the spectral analysis of the entire data. A better agreement between these histograms cannot be expected because the internal waves are strongly dependent on the changing local mean stratification,⁵ whereas the swell is governed by quite different sources.




Figure 4. Histograms of periods of swell modulations (left) and of internal waves (right) at the NEL Tower.

CONCLUSIONS

1. Resonant interaction between amplitude-modulated swell and internal waves may create internal waves of 2-to-20-minute periods. Internal waves with amplitudes of 1-to-3 meters can be produced within an interaction time of 15 minutes if the stratification of the water is such that

$$|\vec{k}_1 - \vec{k}_2|^2 / (\omega_1 - \omega_2)^2$$

is an eigenvalue of the internal wave equation (\vec{k}_1 , \vec{k}_2 , ω_1 , and ω_2 being wave numbers and frequencies of the modulated swell).

2. Internal waves due to amplitude-modulated swell have the same characteristics as the modulation. Specifically, they have the same wavelength and period and travel in the same direction as the amplitude modulation.

3. The resonance process is most efficient in the case of a modulation which travels in the same direction as the carrier wave (the main constituent) of a swell. The creation of internal waves of this type is strongly dependent on the stratification of the water.

4. Preliminary evaluations of temperature and swell records from the NEL Tower show good agreement between periods of swell modulation and internal wave periods.

5. It is likely that internal waves are created on the entire continental shelf off Southern California during times of favorable stratification. This depends on tide and wind.

RECOMMENDATIONS

1. Verify, with additional measurements at the NEL Tower, the theory that internal waves are produced by swell.

2. Acquire, for complete analysis, long records of temperature fluctuations and surface waves in order to compute spectra with high resolution and statistical confidence.

3. Check, by measurements of surface waves at two positions, and slick observations, the result that internal waves travel in the direction of the swell modulation.

REFERENCES

1. Navy Electronics Laboratory Report 937, Slicks and Temperature Structure in the Sea, by E. C. LaFond, 2 November 1959.
2. LaFond, E.C., "Internal Waves," p.731-751 in M.N. Hill, The Sea, v.1, Interscience, 1962.
3. Navy Electronics Laboratory Report 1342, The U. S. Navy Electronics Laboratory's Oceanographic Research Tower; Its Development and Utilization, by E.C. LaFond, 22 December 1965
4. Lee, O.S., "Observations on Internal Waves in Shallow Water," Limnology and Oceanography, v.6, p.312-321, July 1961
5. Cairns, J.L. and LaFond, E.C., "Periodic Motions of the Seasonal Thermocline Along the Southern California Coast," Journal of Geophysical Research, v.71, p.3903-3915, 15 August 1966
6. Ufford, C.W., "Internal Waves Measured at Three Stations," American Geophysical Union. Transactions, v.28, p.87-95, February 1947
7. Ewing, G., "Slicks, Surface Films and Internal Waves," Journal of Marine Research, v.9, p.161-187, 1950

8. Cox, G.S., "Internal Waves, Part 2," p.752-763 in M.N. Hill, The Sea, v.1, Interscience, 1962
9. Ball, F.K., "Energy Transfer Between External and Internal Gravity Waves," Journal of Fluid Mechanics, v.19, p.465-478, July 1964
10. Thorpe, S.A., "On Wave Interactions in a Stratified Fluid," Journal of Fluid Mechanics, v.24, p.737-751, April 1966
11. Kenyon, K.E., Wave-Wave Scattering for Gravity Waves and Rossby Waves, (Ph.D. Thesis, University of California at San Diego), 1966
12. Hasselmann, K., "Feynman Diagrams and Interaction Rules of Wave-Wave Scattering Processes," Reviews of Geophysics, v.4, p.1-32, February 1966
13. Krauss, W., Interne Wellen, v.2; Methoden und Ergebnisse der Theoretischen Ozeanographie, Gebrüder Borntraeger, 1966

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Navy Electronics Laboratory San Diego, California 92152		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE INTERACTION BETWEEN SURFACE AND INTERNAL WAVES IN SHALLOW WATER			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Research and Development Report, August 1966 - December 1966			
5. AUTHOR(S) (First name, middle initial, last name) W. Krauss			
6. REPORT DATE 30 January 1967		7a. TOTAL NO. OF PAGES 28	7b. NO. OF REFS 13
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) 1432	
b. PROJECT NO. SR 104 03 01 Task 0594			
c. (NEL L41171)		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Naval Ship Systems Command Department of the Navy	
13. ABSTRACT <p>The relationship of internal waves, with 2-to-20-minute periods, to surface waves was investigated. It was found that (1) the interaction between amplitude-modulated swell and internal waves may create internal waves of 2-to-20-minute periods; (2) internal waves due to amplitude-modulated swell have the same characteristics as the modulation; and (3) the resonance process is most efficient in the case of a modulation traveling in the same direction as the carrier wave of a swell.</p>			

DD FORM 1473 (PAGE 1)

1 NOV 65

S/N 0101-807-6801

UNCLASSIFIED

Security Classification

<p>Nav Electronics Lab., San Diego, Calif. Report 1432</p> <p>INTERACTION BETWEEN SURFACE AND INTERNAL WAVES IN SHALLOW WATER by W. Krauss, 28 p., 30 Jan 67.</p> <p>UNCLASSIFIED</p> <p>The relationship of internal waves, with 2-to-20-minute periods, to surface waves was investigated. It was found that (1) the interaction between amplitude-modulated swell and internal waves may create internal waves of 2-to-20-minute periods; (2) internal waves due to amplitude-modulated swell have the same characteristics as the modulation; and (3) the resonance process is most efficient in the case of a modulation traveling in the same direction as the carrier wave of a swell.</p>	<ol style="list-style-type: none"> 1. Internal Waves - Shallow Water 2. Ocean Waves - Analysis <p>I. Krauss, W.</p> <p>SR 104 03 01, Task 0594 (INEL L41171)</p> <p>This card is UNCLASSIFIED</p>
<p>Nav Electronics Lab., San Diego, Calif. Report 1432</p> <p>INTERACTION BETWEEN SURFACE AND INTERNAL WAVES IN SHALLOW WATER by W. Krauss, 28 p., 30 Jan 67.</p> <p>UNCLASSIFIED</p> <p>The relationship of internal waves, with 2-to-20-minute periods, to surface waves was investigated. It was found that (1) the interaction between amplitude-modulated swell and internal waves may create internal waves of 2-to-20-minute periods; (2) internal waves due to amplitude-modulated swell have the same characteristics as the modulation; and (3) the resonance process is most efficient in the case of a modulation traveling in the same direction as the carrier wave of a swell.</p>	<ol style="list-style-type: none"> 1. Internal Waves - Shallow Water 2. Ocean Waves - Analysis <p>I. Krauss, W.</p> <p>SR 104 03 01, Task 0594 (INEL L41171)</p> <p>This card is UNCLASSIFIED</p>
<p>Nav Electronics Lab., San Diego, Calif. Report 1432</p> <p>INTERACTION BETWEEN SURFACE AND INTERNAL WAVES IN SHALLOW WATER by W. Krauss, 28 p., 30 Jan 67.</p> <p>UNCLASSIFIED</p> <p>The relationship of internal waves, with 2-to-20-minute periods, to surface waves was investigated. It was found that (1) the interaction between amplitude-modulated swell and internal waves may create internal waves of 2-to-20-minute periods; (2) internal waves due to amplitude-modulated swell have the same characteristics as the modulation; and (3) the resonance process is most efficient in the case of a modulation traveling in the same direction as the carrier wave of a swell.</p>	<ol style="list-style-type: none"> 1. Internal Waves - Shallow Water 2. Ocean Waves - Analysis <p>I. Krauss, W.</p> <p>SR 104 03 01, Task 0594 (INEL L41171)</p> <p>This card is UNCLASSIFIED</p>

INITIAL DISTRIBUTION LIST

CHIEF OF NAVAL MATERIAL
MAT 0331
COMMANDER, NAVAL SHIP SYSTEMS COMMAND
SHIPS 1610
SHIPS 1631
SHIPS 2021 (2)
SHIPS 204113
COMMANDER, NAVAL AIR SYSTEMS COMMAND
AIR 5330
AIR 5401
AIR 604
COMMANDER, NAVAL ORDNANCE SYSTEMS COMMAND
ORD 03C
ORD 0322
ORD 9132
COMMANDER, NAVAL FACILITIES ENGINEERING
COMMAND
FAC 42310
COMMANDER, NAVAL ELECTRONIC SYSTEMS COMMAND
TECHNICAL LIBRARY
COMMANDER, NAVAL SHIP ENGINEERING CENTER
CODE 6120
CODE 6179B
CODE 6179C03
CODE 6360
CHIEF OF NAVAL PERSONNEL
PERS 118
CHIEF OF NAVAL OPERATIONS
OP-03EG
OP-311
OP-312F
OP-322C
OP-07T
OP-702C
OP-71
OP-716
OP-09B5
OP-922Y4C1
CHIEF OF NAVAL RESEARCH
CODE 416
CODE 418
CODE 427
CODE 466
CODE 468
COMMANDER IN CHIEF
US PACIFIC FLEET
CODE 93
US ATLANTIC FLEET
COMMANDER OPERATIONAL TEST AND EVALUATION
FORCE
KEY TEST TEST AND EVALUATION DETACHMENT
DEPUTY COMMANDER OPERATIONAL TEST AND
EVALUATION FORCE, PACIFIC
COMMANDER SUBMARINE FORCE
US PACIFIC FLEET
CODE 21
US ATLANTIC FLEET
COMMANDER ANTISUBMARINE WARFARE FORCE
US PACIFIC FLEET
COMMANDER FIRST FLEET
COMMANDER SECOND FLEET
COMMANDER TRAINING COMMAND
US ATLANTIC FLEET
OFFICE OF THE OCEANOGRAPHER OF THE NAVY
COMMANDER OCEANOGRAPHIC SYSTEM PACIFIC
COMMANDER SUBMARINE DEVELOPMENT GROUP TWO
COMMANDER, DESTROYER DEVELOPMENT GROUP,
PACIFIC
COMMANDER FLEET AIR WINGS, ATLANTIC FLEET
NAVAL AIR DEVELOPMENT CENTER
LIBRARY
NAVAL MISSILE CENTER
TECHNICAL LIBRARY
PACIFIC MISSILE RANGE
CODE 3250
NAVAL ORDNANCE TEST STATION
CHINA LAKE
CODE 753
PASADENA ANNEX
LIBRARY
NAVAL WEAPONS LABORATORY
KXL
PEARL HARBOR NAVAL SHIPYARD
CODE 246P
PORTSMOUTH NAVAL SHIPYARD
CODE 242L
PUGET SOUND NAVAL SHIPYARD
CODE 246
SAN FRANCISCO NAVAL SHIPYARD
HUNTERS POINT DIVISION
NAVAL RADIOLOGICAL DEFENSE LABORATORY
CODE 222A
DAVID TAYLOR MODEL BASIN
LIBRARY
NAVY MINE DEFENSE LABORATORY
CODE 716
NAVAL TRAINING DEVICE CENTER
TECHNICAL LIBRARY

NAVY UNDERWATER SOUND LABORATORY
LIBRARY
CODE 905
ATLANTIC FLEET ASW TACTICAL SCHOOL
LIBRARY
NAVY MARINE ENGINEERING LABORATORY
CODE 257
NAVAL CIVIL ENGINEERING LABORATORY
LS4
NAVAL RESEARCH LABORATORY
CODE 2027
CODE 4320
CODE 5440
NAVAL ORDNANCE LABORATORY
CORONA
TECHNICAL LIBRARY
SILVER SPRING, MD.
DIVISION 221
DIVISION 730
NAVY UNDERWATER SOUND REFERENCE LABORATORY
LIBRARY
FLEET ASW SCHOOL
TACTICAL LIBRARY
FLEET SONAR SCHOOL
NAVAL UNDERWATER WEAPONS RESEARCH AND
ENGINEERING STATION
LIBRARY
OFFICE OF NAVAL RESEARCH BRANCH OFFICE
PASADENA
CHIEF SCIENTIST
BOSTON
CHICAGO
SAN FRANCISCO
LONDON
NAVAL SHIP MISSILE SYSTEMS ENGINEERING
STATION
CODE 903
CHIEF OF NAVAL AIR TRAINING
TRAINING RESEARCH DEPARTMENT
NAVY WEATHER RESEARCH FACILITY
NAVAL OCEANOGRAPHIC OFFICE
CODE 1640
SUPERVISOR OF SHIPBUILDING, US NAVY
GROTON, CONN.
CODE 249
NAVAL POSTGRADUATE SCHOOL
DEPT. OF ENVIRONMENTAL SCIENCES
LIBRARY
FLEET NUMERICAL WEATHER FACILITY
NAVAL APPLIED SCIENCE LABORATORY
CODE 9200
CODE 9830
NAVAL ACADEMY
ASSISTANT SECRETARY OF THE NAVY
(RESEARCH AND DEVELOPMENT)
NAVAL SECURITY GROUP
C-43
AIR DEVELOPMENT SQUADRON ONE
VX-1
SUBMARINE FLOTILLA ONE, US PACIFIC FLEET
DEFENSE DOCUMENTATION CENTER (20)
DEPARTMENT OF DEFENSE RESEARCH AND
ENGINEERING
WEAPONS SYSTEMS EVALUATION GROUP
DEFENSE ATOMIC SUPPORT AGENCY
DOCUMENT LIBRARY SECTION
NATIONAL OCEANOGRAPHIC DATA CENTER
CODE 2400
COAST GUARD OCEANOGRAPHIC UNIT
NATIONAL ACADEMY OF SCIENCES/
NATIONAL RESEARCH COUNCIL
COMMITTEE ON UNDERSEA WARFARE
COAST GUARD HEADQUARTERS
OSR-2
ARCTIC RESEARCH LABORATORY
WOODS HOLE OCEANOGRAPHIC INSTITUTION
DOCUMENT LIBRARY LO-206
ENVIRONMENTAL SCIENCE SERVICE ADM.
COAST AND GEODETIC SURVEY
ROCKVILLE, MD.
WASHINGTON SCIENCE CENTER - 23
WASHINGTON, D. C.
US WEATHER BUREAU
DIRECTOR, METEOROLOGICAL RESEARCH
LIBRARY
GEOLOGICAL SURVEY LIBRARY
DENVER SECTION
ESSA/INSTITUTE FOR TELECOMMUNICATION
SCIENCES AND AERONOMY
BOULDER, COLO.
FEDERAL COMMUNICATIONS COMMISSION
RESEARCH DIVISION
NATIONAL SEVERE STORMS LABORATORY
CENTRAL INTELLIGENCE AGENCY
OCR/DD-STANDARD DISTRIBUTION
NATIONAL BUREAU OF STANDARDS
BOULDER, COLO.

BUREAU OF COMMERCIAL FISHERIES
LA JOLLA, CALIF.
TUNA RESOURCES LABORATORY LA JOLLA
WASHINGTON, D. C.
BRANCH OF MARINE FISHERIES
WOODS HOLE, MASS.
BIOLOGICAL LABORATORY LIBRARY
HONOLULU, HAWAII
FISH AND WILDLIFE SERVICE LIBRARY
STANFORD, CALIF.
BIOLOGICAL LABORATORY
ABERDEEN PROVING GROUND
TECHNICAL LIBRARY
ARMY MISSILE CENTER
REDSTONE SCIENTIFIC INFORMATION CENTER
DOCUMENT SECTION
ARMY ELECTRONICS RESEARCH AND DEVELOPMENT
LABORATORY
ARMY ELECTRONICS COMMAND
MANAGEMENT & ADMINISTRATIVE SERVICES DEPT
AMSEL-RD-MAT
COASTAL ENGINEERING RESEARCH CENTER
ARMY CORPS OF ENGINEERS
AIR FORCE HEADQUARTERS
DIRECTOR OF SCIENCE AND TECHNOLOGY
ARST-A
AIR UNIVERSITY LIBRARY
AULT3-5028
AIR FORCE EASTERN TEST RANGE
AFMTC TECHNICAL LIBRARY - MU-135
AIR PROVING GROUND CENTER
PGP5-12
HEADQUARTERS AIR WEATHER SERVICE
AW555/5IPD
WRIGHT-PATTERSON AIR FORCE BASE (1)
SYSTEMS ENGINEERING GROUP (RTD)
SEPIR
UNIVERSITY OF MICHIGAN
OFFICE OF RESEARCH ADMINISTRATION
NORTH CAMPUS
COOLEY ELECTRONICS LABORATORY
UNIVERSITY OF CALIFORNIA-SAN DIEGO
MARINE PHYSICAL LABORATORY
SCRIPPS INSTITUTION OF OCEANOGRAPHY
LIBRARY
UNIVERSITY OF MIAMI
THE MARINE LABORATORY LIBRARY
MICHIGAN STATE UNIVERSITY
LIBRARY-DOCUMENTS DEPARTMENT
COLUMBIA UNIVERSITY
LAMONT GEOLOGICAL OBSERVATORY
DARTMOUTH COLLEGE
RADIOPHYSICS LABORATORY
CALIFORNIA INSTITUTE OF TECHNOLOGY
JET PROPULSION LABORATORY
HARVARD COLLEGE OBSERVATORY
HARVARD UNIVERSITY
GORDON MCKAY LIBRARY
LYMAN LABORATORY
OREGON STATE UNIVERSITY
DEPARTMENT OF OCEANOGRAPHY
UNIVERSITY OF WASHINGTON
DEPARTMENT OF OCEANOGRAPHY
FISHERIES-OCEANOGRAPHY LIBRARY
APPLIED PHYSICS LABORATORY
NEW YORK UNIVERSITY
DEPARTMENT OF METEOROLOGY AND OCEANOGRAPHY
UNIVERSITY OF ALASKA
GEOGRAPHICAL INSTITUTE
UNIVERSITY OF RHODE ISLAND
NARRAGANSETT MARINE LABORATORY
LIBRARY
YALE UNIVERSITY
BINGHAM OCEANOGRAPHIC LABORATORY
FLORIDA STATE UNIVERSITY
OCEANOGRAPHIC INSTITUTE
UNIVERSITY OF HAWAII
HAWAII INSTITUTE OF GEOPHYSICS
ELECTRICAL ENGINEERING DEPARTMENT
AGM COLLEGE OF TEXAS
DEPARTMENT OF OCEANOGRAPHY
THE UNIVERSITY OF TEXAS
DEFENSE RESEARCH LABORATORY
ELECTRICAL ENGINEERING RESEARCH LABORATORY
PENNSYLVANIA STATE UNIVERSITY
ORDNANCE RESEARCH LABORATORY
STANFORD RESEARCH INSTITUTE
NAVAL WARFARE RESEARCH CENTER
MASSACHUSETTS INSTITUTE OF TECHNOLOGY
ENGINEERING LIBRARY
LINCOLN LABORATORY
RADIO PHYSICS DIVISION
FLORIDA ATLANTIC UNIVERSITY
DEPARTMENT OF OCEAN ENGINEERING
THE JOHNS HOPKINS UNIVERSITY
APPLIED PHYSICS LABORATORY
DOCUMENT LIBRARY
INSTITUTE FOR DEFENSE ANALYSES
DOCUMENT LIBRARY